



TRAUMATIC PELVIC RING INJURIES

POPULATION



Patients admitted to
participating hospitals
during study period

OBSERVATIONS



Demographic parameters
(comorbidities, pre-injury
functional status, trauma
mechanism, fracture
classification)



Diagnostic imaging used
(X-ray, CT, MRI)



Emergent surgical
intervention &
complications



Definitive surgical or
conservative fracture
treatment & complications

PATHOLOGY



Traumatic
Pelvic
Ring
Injuries

METHODS

- Prospective 90-day observational cohort study, with 180-day follow-up
- Validated methodology
- REDCap database (secure)
- Authorship and data-ownership for collaborators

OUTCOME MEASURES



Overall and injury-related mortality



Clinical outcome (pain, neur-
ological-urological-functional
deficiencies)



Radiological outcome
(osseous healing, residual
posterior and/or anterior
displacement)



Social reintegration



Surgical complications requiring re-
operation

STEERING GROUP

Bass G. (USA), Mohseni S. (SWE), Bolierakis E. (GER), Höch A. (GER),
Schipper I.B. (NL), Pape H.-C. (CH), Hildebrand F. (GER).*(chair)

(Information: Efthychios Bolierakis, University Hospital RWTH Aachen,
ebolierakis@ukaachen.de)